

ARTICLE

Anatomy of positive messages in healthcare consultations: component analysis of messages within 22 randomised trials

Jeremy Howick PhD^a, Emily Lyness MB ChB^b, Charlotte Albury DPhil^c, Kirsten A. Smith PhD^d, Hajira Dambha-Miller MRCP^e, Mohana Ratnapalan MBBS^f, Jane Vennik PhD^g, Stephanie Hughes MSc^h, Jennifer Bostockⁱ, Leanne Morrison PhD^j, Christian Mallen PhD^k, Hazel Everitt MB ChB BSc MSc PhD FRCGP^l, Sue Dean RN BA MA^m, Tracy Levett-Jones PhDⁿ, Serra Ivynian PhD^o, Paul Little MBBS BA MD MRCP FRCGP FMedSci^p and Felicity L. Bishop PhD^q

- a Director, Oxford Empathy Programme & Senior Researcher and Impact Fellow, Faculty of Philosophy, University of Oxford, Radcliffe Observatory Quarter, Oxford, UK
- b Academic Clinical Fellow with the School of Primary Care, Population Sciences and Medical Education, University of Southampton, Southampton, UK
- c Research Fellow, Department of Primary Care Health Sciences, University of Oxford, Oxford, UK
- d Research Fellow, Department of Psychology, University of Southampton, Southampton, UK
- e Clinical Lecturer, Academic Unit of Primary Care & Population Science, University of Southampton, Southampton, UK
- f Academic Clinical Fellow, School of Primary Care, Population Sciences and Medical Education, Faculty of Medicine, University of Southampton, Southampton, UK
- g Research Fellow, Primary Medical Care, University of Southampton, Southampton, UK
- h Senior Research Assistant & Postgraduate Student, Academic Unit of Primary Care & Population Science, University of Southampton, Southampton, UK
- i Research Ethics and PPI Advisor, Department of Psychological Medicine, King's College London, London, UK
- j Lecturer in Health Psychology, Centre for Community and Clinical Applications of Health Psychology, Faculty of Environmental and Life Sciences and the School of Primary Care, Population Sciences and Medical Education, Faculty of Medicine, University of Southampton, Southampton, UK
- k Director & NIHR Professor of General Practice, School for Primary, Community and Social Care, Keele University, Newcastle under Lyme, UK
- l Professor of Primary Care Research, Academic Unit of Primary Care & Population Science, University of Southampton, Southampton, UK
- m Lecturer, Faculty of Health, University of Technology Sydney, Ultimo, NSW, Australia
- n Professor of Nursing Education, Faculty of Health, University of Technology Sydney, Ultimo, NSW, Australia
- o Research Fellow, Nursing, Faculty of Health, University of Technology Sydney, Ultimo, NSW, Australia
- p Professor of Primary Care Research, Academic Unit of Primary Care & Population Science, University of Southampton, Southampton, UK
- q Associate Professor & Lecturer in Health Psychology, Faculty of Social & Human Sciences, University of Southampton, Southampton, UK

Abstract

Background: Patients desire both honesty and hope from their healthcare practitioners. A recent systematic review found that healthcare practitioners who deliver positive messages improve patient outcomes, most notably by reducing pain. However, the verbal and non-verbal components of positive messages within these trials varied greatly, which presents a barrier to the implementation of this evidence and person-centered care.

Objective: To investigate common components of positive messages within the reviewed trials.

Methods: We extracted the verbal and non-verbal language used to deliver positive messages in 22 trials from a recent systematic review. Three independent researchers coded the components of the messages using content analysis.

Results: Positive messages in our sample had between 2 and 18 different components. These were clustered into 5 areas: specifying the positive outcomes, making the message personal, drawing on associations and meanings, providing a supportive psychological context and providing a rationale. Messages were reinforced through repetition in half the studies. Within the clusters, the most common components of positive messages were suggestions of specific effects (18 studies) and personalised formulations (15 studies). Most studies did not describe the components of positive messages adequately.

Conclusions: Positive messages within randomized trials are complex interventions, with most including strong suggestions about specific effects, presented confidently and made personally relevant to the individual patient. Future trials of positive messages should report all components of these interventions.

Keywords

Communication, communication skills, complex interventions, empathy, ethics, mechanisms of action, pain, pain reduction, person-centered healthcare, positive communication, randomised controlled trials

Correspondence address

Dr. Jeremy Howick, Faculty of Philosophy, University of Oxford, Radcliffe Infirmary Quarter, Woodstock Road, Oxford, OX2 6GF, UK. E-mail: Jeremy.howick@philosophy.ox.ac.uk

Accepted for publication: 31 December 2019

Introduction

Communication between patients and practitioners lies at the heart of person-centered healthcare, and good communication has many potential benefits including improved patient satisfaction [1-3], reduced medico-legal risk [4], increased patient safety [1,5] and improved adherence to medication [6]. Some evidence also suggests that enhanced communication may reduce healthcare practitioner burnout [7,8]. One aspect of patient-practitioner communication that is emerging as having a strong evidence base is the communication to patients of positive messages (encouraging the patient to believe they will experience a positive outcome) [9,10]. Encouraging positive patient expectations can lead to better outcomes [11-13] and even patients with serious illnesses desire (realistic) hope [14]. In addition, positive messages are taken to be a core component of empathy [15], which, in turn, is a core component of person-centered care [16].

Mechanisms by which positive messages may improve patient outcomes have been investigated at psychological and neurophysiological levels. The practitioner's positive message is assumed to lead to the patient's expectation of a positive outcome. In turn, the patients' outcome expectancies [17,18] are positively associated with actual symptom reduction [17,19]. At a neurophysiological level, positive outcome expectancies are thought to activate the neuronal reward mechanisms including increased dopamine activity in the nucleus accumbens [20,21].

The potential benefits of positive messages have recently been quantified in a systematic review of 22 randomised trials. The review included randomised trials that measured clinical outcomes and excluded other types of studies (such as observational studies), as well as trials that did not include clinical outcomes (such as experimentally induced pain). The trials included conditions ranging from chronic pain to coronary artery bypass surgery. Positive messages were found (by a small to modest amount) to reduce patient pain, anxiety, morphine use, length of stay in hospital, and improve lung function [22].

While the systematic review demonstrated that delivering positive messages can improve patient outcomes, a barrier to implementing the evidence is heterogeneity. The messages delivered to patients ranged from very simple sentences, (e.g., "I think this will work for you") to more comprehensive messages encouraging a positive outlook [23] and giving clear diagnoses [24]. The types of practitioners, training given to practitioners,


medical conditions, and type of benefits to patients (such as reduced pain), also varied within the trials [22]. The heterogeneity makes it difficult to infer what healthcare practitioners should say or do in order to deliver an effective positive message. The aim of this study was to identify and describe the components of effective positive messages tested in the trials from a recently published systematic review [22].

Methods

We used a conventional content analysis [25,26] to describe the components of positive messages trialled in the systematic review of positive communication. The sections of the 22 studies describing the positive messages (reported across 20 papers) [3,23,24,27-43] were imported into NVivo for analysis. We also examined supplementary online material and one additional paper [44]. Descriptions of the positive messages were taken to include: the content and context of the positive messages as described by the authors, direct scripts or images showing the positive message (where available) and descriptions of non-verbal aspects of the message (tone of voice, body language, attire), as well as details of any other manipulation that might convey or support a positive message.

We then coded this material inductively using labels that summarised individual components of the positive message. Initial codes closely reflected the original authors' own terminology and concepts and were not mutually exclusive. We then compared and contrasted initial codes and their exemplars across the studies, merging them into categories and sub-categories based on common features that transcended idiosyncratic study-specific descriptions. All categories were then defined, and a coding manual was produced capturing category names, definitions, and examples. We used our knowledge of multiple theories (including response expectancy theory, social cognitive theory, common sense model of illness representations, theories of persuasive communication and attitude change) to cluster the categories according to what these components and features were doing. For example, some categories described components that directly conveyed information about the likely outcomes of a treatment (and were clustered as 'specifying the positive outcome(s)'), while other categories described components that explained to patients why the clinician believed the intervention would have positive effects (and were clustered as 'providing a rationale'). To refine the coding

Figure 1 Overview of Components of Positive Messages



manual and ensure the coding process could be reliably replicated, two authors read and independently coded half of the papers each using the coding manual. This process identified some discrepancies which were discussed and resolved by discussion and, where required, adjustment of the coding manual. We used NVivo queries to help us examine patterns in the data and count the number of distinct components used in each study.

Results

We identified 25 distinct components of positive messages (see Figure 1). Included studies used between 2 [33] and 18 [3] different components to convey their positive message (Median = 4; Mean = 5.8). We clustered these components into 5 areas (see Table 1 for details):

1. Specifying the positive outcome(s).
2. Making the message personal and accessible.
3. Conveying a positive message through enhancing associations.
4. Encouraging a supportive psychological context.
5. Providing a rationale.

Eleven studies (50%) reinforced their positive message by presenting it repeatedly. Table 2 specifies which studies included components from each of the 5 clusters.

Table 1 Overview of Components of Positive Messages

Cluster	Component	Studies reporting component (n)
Specifying the positive outcome(s)	Positive suggestion of specific effects	18
	Strong or definite assertion of positive outcome	14
	Shaping expectations about immediacy of effects	10
	Positive suggestion emphasised through adjectives	10
	Cautious or probabilistic assertion of positive outcome	8
	Positive suggestion of general effects	7
	Shaping expectations about duration of effects	2
Making the message personal and accessible	Message formulation is direct and personal	15
	Positive suggestion conveyed through analogy	2
Conveying positive message through associations and meanings	Increasing attention to administration	9
	Sensory properties of medication	4
	Social modelling or peer identification	4
	Meaning of dosage	2
	Communicating authority	2
	Branding	1
Encouraging a supportive psychological context	Supporting helpful symptom interpretation	8
	Empathy and respect for patient	3
	Supporting patient's sense of autonomy	3
	Enhancing self-efficacy	2
	Allaying safety concerns	2
Appeal to evidence Providing a rationale	Message evidenced by verbal appeals to other patients	7
	Rationale for how the intervention may result in positive outcomes	6
	Message evidenced by verbal appeals to evidence	3
	Message evidenced by verbal appeals to clinician's experience	2
	Message evidenced by verbal appeals to institutional authority	1

Table 2 Component Clusters Identified in each Study

Study	Specifying the positive outcome(s)	Making the message personal and accessible	Conveying positive message through associations and meanings	Encouraging a supportive psychological context	Providing a rationale (including appeal to experience and authority)
1. Benedetti 2003 [46]	✓	✓	✓	✗	✗
2. Benedetti 2003 [64]	✓	✓	✓	✗	✗
3. Benedetti 2003 [65]	✓	✗	✓	✗	✗
4. deCraen 2001 [28]	✓	✗	✓	✗	✓
5. Dutt-Gupta 2007 [29]	✓	✗	✗	✓	✓
6. Goodenough 1997 [30]	✓	✓	✓	✗	✗
7. Kemeny 2007 [31]	✓	✓	✓	✓	✗
8. Knipschild 2005 [32]	✓	✓	✗	✓	✗
9. Lauder 1995 [33]	✓	✗	✗	✗	✗
10. Olsson 1989 [34]	✓	✓	✗	✓	✗
11. Petersen 2014 [35]	✓	✗	✓	✗	✓
12. Petersen 2012 [36]	✓	✗	✓	✗	✓
13. Phillips 2006 [37]	✓	✓	✓	✓	✓
14. Resnick 1996 [38]	✓	✓	✓	✓	✓
15. Rief 2017 [23]	✓	✗	✗	✓	✗
16. Ronel 2011 [39]	✓	✓	✓	✗	✓
17. Suarez 2010 [40]	✓	✓	✗	✗	✓
18. Szilagyi 2007 [43,44]	✓	✓	✓	✓	✓

19. Thomas 1987 [24]	✓	✓	✓	✓	✗
20. Varelmann 2010 [41]	✓	✓	✓	✗	✗
21. Wang 2008 [42]	✓	✓	✗	✗	✓
22. Wise 2009 [3]	✓	✓	✓	✓	✓

Specifying the positive outcome(s)

The messages varied in how specific the outcome was anticipated to be. Eighteen studies (82%) made a positive suggestion of specific effects such as “[your] pain will subside in a few minutes” [27]. In fewer cases (7, 32%) the messages was more general, for example, about “getting better,” or the intervention being “effective,” or communicating that the patient would be “living active and fulfilling lives” [3]. Ten studies (45%) specified how quickly the patient could expect to benefit and typically provided for a little variation within these descriptions, for example, “[This drug] will decrease the pain quickly after taking it” [28] and “You will be better within a week or so” [32]. Two studies (9%) specified how long the benefit should last. In 14 studies (64%), outcomes were specified using strong assertions such as telling patients “their pain was going to subside within a few minutes” [45] or that “the treatment would certainly make [the patient] better” [24]. Fewer studies (8, 36%) specified positive outcomes using more cautious or probabilistic language. For example, in a study of positive messaging for venepuncture in children, practitioners were instructed to say “I am going to put some cream on your arm that *might* make it (the needle) hurt less” [30]. Ten studies (45%) used adjectives to emphasise a likely positive outcome in relation to analgesics/anaesthetics, painful procedures, and asthma; examples include “this drug is very effective” [39] and “we are trying out a new special cream” [30]. Stronger adjectives alluding to the power of drugs, such as “potent” [45] were also used to describe analgesics.

Making the message personal and accessible

Fifteen studies (68%) made positive messages personal and direct (person-centered) by referencing the patient’s name, directing the message to them personally as in “I think this will work for you” [40] or writing a personalised handwritten prescription or instruction. Two studies (9%) used analogy to make the message accessible and engaging to patients. For example, in a study of positive suggestion for ventilated patients: “Some imagine that these noises [made by the machines in intensive care] are that of a pleasant boat-ride, a cruise” [43,44].

Conveying a positive message through enhancing associations and meanings

Sometimes positive messages included references to positive associations or meanings patients might be likely to have. The most common of these (reported by 9 studies) involved increasing the patient’s attention to treatment.

This was achieved by, for example, encouraging the patient to observe an injection being given. Presumably the patients were assumed to associate a positive response with the medical procedure [46]. Messages in 4 studies aimed to call the patients’ attention to the sensory properties of medication (such as taste, colour or texture). These properties have been shown in some studies to modify effects [47]. For example, in one study the practitioners were instructed to say, “We are going to inject the local anaesthetic that will numb the area where we are going to do the epidural/spinal anaesthesia and you will be comfortable during the procedure” [41]. In 4 studies, the patient was compared with other similar patients. For example, one study stated: “The commercial included positive messages showing attractive young adults with asthma leading apparently active healthy and fulfilling lives” [3]. This was intended to help the patient associate themselves with these healthy, fulfilled people in the commercials. Communicating authority also featured in 2 of the messages. This was accomplished in some trials by training practitioners to speak with authority and conviction [31], by modifying their appearance (such as having them wear a white coat) [36] or through the mere presence of a qualified, authoritative individual (a doctor as opposed to a technician). Interventions were branded in one study, in order (according to the study) to benefit from the association between branding and increased perceived potency [3].

Encouraging a supportive psychological context

Eight studies (36%), provided a supportive psychological context. This was achieved by validating patients’ experiences, reassuring patients about their symptoms, guiding patients about likely symptom trajectories for recovery, and helping to normalise the side effects of an intervention. For example, practitioners in one study were encouraged to “explain that what they are experiencing may not be an uncommon symptom considering what they have been enduring with their debilitation and hospitalization” [37]. Three studies included components that communicated empathy and respect to the patient, for example, by having the healthcare practitioner introduce themselves, shaking hands, and/or exploring the patients’ perspective and emotional wellbeing. For instance, in one study of self-efficacy and outcome expectations in rehabilitation, patients are encouraged to identify their thoughts and feelings [38]. Three studies supported patient autonomy, by encouraging personal goal setting through the use of mental imagery and personal ideation. In one study of pre-operative expectations and recovery from cardiac surgery “patients were encouraged to develop

personal ideas and images about their future after surgery, including plans about activities and how they will enjoy their life afterwards (outcome expectations)” [23]. Two studies, both in rehabilitation settings, included components that enhanced patients’ self-efficacy alongside their outcome expectations. For example, rehabilitation patients were encouraged to be realistically optimistic about their capacity to reach a goal, to find examples of previous success in order to engender a sense of mastery and to understand that the healthcare team believed patients were capable of achieving their stated goals [38]. Two studies used supportive comments to reassure the patient of the safety of the intervention. Practitioners in a study of acute care were instructed to say: “The machines around you continuously monitor the needs of your body. The beeps are signals for us, just serving your safety” [43,44].

Providing a rationale (including appeal to evidence and experience)

There were two types of rationales provided in some of the studies. The first described the mechanism of action of the intervention. For example, a study in rehabilitation noted that the exercises would lead to improved strength, leading to improved confidence, and therefore improved function [37]. Another study stated that the experimental drug “improves many aspects of asthma by blocking leukotrienes” [3]. The second type of rationale was a reason why the practitioner believed (and hence why the patient should expect) the intervention to be effective. These rationales included appeals to other patients’ experience (7, 31%), external evidence (3, 14%), clinician experience (2, 10%), and institutional authority (1, 5%). Appeal to other patients’ experience often involved simple messages such as “most of my patients get better” [40]. Three of the studies appealed to evidence (such as “The PCA pump was very effective in removing the postoperative pain affliction” [42]) and one study to an institutional authority (such as “the US Food and Drug Administration has judged it to be safe and effective for asthma” [3]).

Other features of positive messages: reinforcement through repetition

Repetition was a ‘meta feature’ of the positive messages, and was used in 11 (50%) of studies. Repetition occurred before or during treatment administration [3], by multiple individuals (e.g., trial and medical personnel) [33] or in different formats (such as in person and *via* digital or printed material) [37]. The timing of positive message provision varied. Timing of the intervention is another ‘meta feature’ and whether the message was delivered before, during, or after varied, and is potentially important. Most studies (n=14) conveyed their positive message before administering the treatment (typically immediately before administration or prescription). Six studies (27%) conveyed their positive message while administering the treatment (typically for procedural interventions) and 3

(14%) conveyed their positive message after administering the treatment (having already conveyed it prior to administration) [35-37].

Discussion

Positive messages used by healthcare practitioners to improve patient outcomes within randomized clinical trials are often complex interventions that comprise multiple components. Strong verbal suggestions of specific effects are the most common features of positive messages. Other clusters of components that may influence the effectiveness of positive messages include making the message personal and accessible; conveying a positive message through enhancing associations; encouraging a supportive psychological context and providing a rationale.

Our paper adds to the literature on the clinical benefits of positive messages by providing more detail about the content of these messages [10,22,48]. Our study also reflects what many mechanism studies have shown, namely that positive messages can activate the neuronal reward mechanisms including increased dopamine activity in the nucleus accumbens [49] as well as endorphins [50]. In addition, previous literature shows that patients’ pre-existing expectations regarding treatment, influence compliance, attrition and outcomes across a wide range of clinical presentations and subsequent treatments [51-53]. A positive message could - indeed is intended to - encourage patients to have more positive pre-intervention expectations. In addition, some of the components of positive messages that we have identified can be mapped to previously studied sources of outcome expectancies such as (i) repeated pairings of specific stimuli and outcomes (conditioning), (ii) observing others’ responses to stimuli (social learning) and (iii) being persuaded about the likely response to stimuli (instructional learning) [54]. Specifying the positive message also uses instructional learning to generate positive outcome expectancies. Our findings also related to the research on persuasion [55], motivation [56], (improving) illness perception [57,58], empathic care (which includes positive communication) [59], self-determination theory (by using expectations to promote patient autonomy) [60], enhancing self-efficacy [61] and medication adherence [62].

This is the only study we are aware of that analyses the components of positive messages. We achieved this by examining randomised trials that had been identified within a systematic review. This approach allowed us to focus on good quality trials (the systematic review excluded more potentially biased, non-randomised, trials). There were also some limitations to this study. The first is that the interventions often lacked detailed description of the content and delivery of the positive messages. Hence, the messages actually used may have included additional components that we did not identify. More complete descriptions may have generated more components in positive messages and possibly a different distribution. Another limitation is that we do not know which component (or combination of components) was most

effective for generating the clinical effect. This might have been achieved by comparing components of positive messages that were effective with components of positive messages that were not effective (this was not possible using this dataset, as all but one of the 22 studies in our sample review reported positive effects). The fact that a component was frequently used does not necessarily indicate that it is most effective. Mitigating this limitation somewhat, all but one of the trials suggested a positive effect of the package of components, which provides weak support for the claim that the common components were effective. Finally, the ethics of positive communication were not explored. Even if positive messages are effective, it does not imply that they are ethically acceptable to patients as they risk violating patient autonomy if they prove to be deceptive.

Future studies of positive messages should consider and build on this evidence base, describe the components of their interventions adequately, and investigate which components of positive messages are most effective. Pragmatic studies and training with healthcare practitioners could consider the extent to which their current communication of positivity reflects the research evidence.

Conclusion

Positive messages used by healthcare practitioners to improve patient outcomes within randomized clinical trials are often complex. Many include repeated strong suggestions about specific effects, are presented confidently, and made personally relevant to the individual patient. A limitation is that interventions are often not well described. Future trials of positive messages should report all components of their interventions. Our detailed analysis could inform more effective intervention development, assessment, as well as clinical practice.

Acknowledgements and Conflicts of Interest

This study was partly funded by the National Institutes of Health Research School for Primary Care Research Award Number 389 (NIHR SPCR 389). Mohana Ratanapalan is a GP ACF who is partly funded by the National Institute for Health Research (NIHR) School for Primary Care Research. The views expressed are those of the author(s) and not necessarily those of the NIHR, NHS or the Department of Health and Social Care. The authors declare no conflicts of interest.

References

- [1] Vermeir, P., Vandijck, D., Degroote, S., Peleman, R., Verhaeghe, R., Mortier, E., Hallaert, G., Van Daele, S., Buylaert, W. & Vogelaers, D. (2015). Communication in healthcare: a narrative review of the literature and practical recommendations. *International Journal of Clinical Practice* 69 (11) 1257-1267.
- [2] Little, P., White, P., Kelly, J., Everitt, H. & Mercer, S. (2015). Randomised controlled trial of a brief intervention targeting predominantly non-verbal communication in general practice consultations. *British Journal of General Practice* 65 (635) e351-356.
- [3] Wise, R.A., Bartlett, S.J., Brown, E.D., Castro, M., Cohen, R., Holbrook, J.T., Irvin, C.G., Rand, C.S., Sockrider, M.M., Sugar, E.A. & American Lung Association Asthma Clinical Research Centers. (2009). Randomized trial of the effect of drug presentation on asthma outcomes: the American Lung Association Asthma Clinical Research Centers. *Journal of Allergy and Clinical Immunology* 124 (3) 436-444, 44e1-8.
- [4] Moore, P.J., Adler, N.E. & Robertson, P.A. (2000). Medical malpractice: the effect of doctor-patient relations on medical patient perceptions and malpractice intentions. *Western Journal of Medicine* 173 (4) 244-250.
- [5] Doyle, C., Lennox, L. & Bell, D. (2013). A systematic review of evidence on the links between patient experience and clinical safety and effectiveness. *BMJ Open* 3 (1) pii: e001570.
- [6] Zolnieriek, K.B. & Dimatteo, M.R. (2009). Physician communication and patient adherence to treatment: a meta-analysis. *Medical Care* 47 (8) 826-834.
- [7] Rotenstein, L.S., Torre, M., Ramos, M.A., Rosales, R.C., Guille, C., Sen, S. & Mata, D.A. (2018). Prevalence of Burnout Among Physicians: A Systematic Review. *Journal of the American Medical Association* 320 (11) 1131-1150.
- [8] Sturzu, L., Lala, A., Bisch, M., Gutter, M., Dobre, D. & Schwan, R. (2019). Empathy and Burnout - A Cross-Sectional Study Among Mental Healthcare Providers in France. *Journal of Medicine and Life* 12 (1) 21-29.
- [9] Howick, J., Lewith, G., Mebius, A., Fanshawe, T., Bishop, F.L., van Osch, M., van Dulmen, S., Christelis, N., Kaptchuk, T. & Mistiaen, P. (2017). Positive messages may reduce patient pain: A meta-analysis. *European Journal of Integrative Medicine* 11, 31-38.
- [10] Peerdeman, K.J., van Laarhoven, A.I., Keij, S.M., Vase, L., Rovers, M.M., Peters, M.L. & Evers, A.W. (2016). Relieving patients' pain with expectation interventions: a meta-analysis. *Pain* 157 (6) 1179-1191.
- [11] Kelley, J.M., Lembo, A.J., Ablon, J.S., Villanueva, J.J., Conboy, L.A., Levy, R., Marci, C.D., Kerr, C.E., Kirsch, I., Jacobson, E.E., Riess, H. & Kaptchuk, T.J. (2009). Patient and practitioner influences on the placebo effect in irritable bowel syndrome. *Psychosomatic Medicine* 71 (7) 789-797.
- [12] Kaptchuk, T.J., Kelley, J.M., Conboy, L.A., Davis, R.B., Kerr, C.E., Jacobson, E.E., Kirsch, I., Schyner, R.N., Nam, B.H., Nguyen, L.T., Park, M., Rivers, A.L., McManus, C., Kokkotou, E., Drossman, D.A., Goldman, P. & Lembo, A.J. (2008). Components of placebo effect: randomised controlled trial in patients with irritable bowel syndrome. *British Medical Journal* 336 (7651) 999-1003.
- [13] Hojat, M., Louis, D.Z., Markham, F.W., Wender, R., Rabinowitz, C. & Gonnella, J.S. (2011). Physicians' empathy and clinical outcomes for diabetic patients. *Academic Medicine* 86 (3) 359-364.

- [14] Mack, J.W. & Smith, T.J. (2012). Reasons why physicians do not have discussions about poor prognosis, why it matters, and what can be improved. *Journal of Clinical Oncology* 30 (22) 2715-2717.
- [15] Mercer, S.W., Maxwell, M., Heaney, D. & Watt, G.C. (2004). The consultation and relational empathy (CARE) measure: development and preliminary validation and reliability of an empathy-based consultation process measure. *Family Practice* 21 (6) 699-705.
- [16] Hardman, D. & Howick, J. (2019). The friendly relationship between therapeutic empathy and person-centered care. *European Journal for Person Centered Healthcare* 7 (2) 351-357.
- [17] Kirsch, I. (1997). Response expectancy theory and application: A decennial review. *Applied and Preventive Psychology* 6 (2) 69-79.
- [18] Kirsch, I. (2004). Conditioning, expectancy, and the placebo effect: Comment on Stewart-Williams and Podd (2004). *Psychological Bulletin* 130 (2) 341-343.
- [19] Kirsch, I., Kong, J., Sadler, P., Spaeth, R., Cook, A., Kaptchuk, T. & Gollub, R.. Expectancy and Conditioning in Placebo Analgesia: Separate or Connected Processes? *Psychology of Consciousness* 1 (1) 51-59.
- [20] Benedetti, F., Mayberg, H.S., Wager, T.D., Stohler, C.S. & Zubieta, J.K. (2005). Neurobiological mechanisms of the placebo effect. *Journal of Neuroscience* 25 (45) 10390-10402.
- [21] Benedetti, F. (2009). Placebo effects: understanding the mechanisms in health and disease. Oxford: Oxford University Press.
- [22] Howick, J., Moscrop, A., Mebius, A., Fanshawe, T.R., Lewith, G., Bishop, F.L., Mistiaen, P., Roberts, N.W., Dieninytė, E., Hu, X.Y., Aveyard, P. & Onakpoya, I.J. (2018). Effects of empathic and positive communication in healthcare consultations: a systematic review and meta-analysis. *Journal of the Royal Society of Medicine* 111 (7) 240-252.
- [23] Rief, W., Shedden-Mora, M.C., Laferton, J.A., Auer, C., Petrie, K.J., Salzmann, S., Schedlowski, M. & Moosdorf, R. (2017). Preoperative optimization of patient expectations improves long-term outcome in heart surgery patients: results of the randomized controlled PSY-HEART trial. *BMC Medicine* 15 (1) 4.
- [24] Thomas, K.B. (1987). General practice consultations: is there any point in being positive? *British Medical Journal (Clinic Research Edition)* 294 (6581) 1200-1202.
- [25] Hsieh, H.F. & Shannon, S.E. (2005). Three approaches to qualitative content analysis. *Qualitative Health Research* 15 (9) 1277-1288.
- [26] Joffe, H. & Yardley, L. (2004). Content and thematic analysis. In: Research methods for clinical and health psychology. Marks, D.F., (Ed.), pp. 56-68. London: Sage.
- [27] Benedetti, F., Maggi, G., Lopiano, L., Lanotte, M., Rainero, I., Vighetti, S. & Pollo, A. (2003). Open versus hidden medical treatments: The patient's knowledge about a therapy affects the therapy outcome. *Prevention & Treatment* 2003; 6 (1) 1a.
- [28] de Craen, A.J.M., Lampe-Schoenmaeckers, A.J.E.M., Kraal, J.W., Tijssen, J.G.P. & Kleijnen, J. (2001). Impact of Experimentally-Induced Expectancy on the Analgesic Efficacy of Tramadol in Chronic Pain Patients: A 2 + 2 Factorial, Randomized, Placebo-Controlled, Double-Blind Trial. *Journal of Pain and Symptom Management* 21 (3) 210-217.
- [29] Dutt-Gupta, J., Bown, T. & Cyna, A.M. (2007). Effect of communication on pain during intravenous cannulation: a randomized controlled trial. *British Journal of Anaesthesia* 99 (6) 871-875.
- [30] Goodenough, B., Kampel, L., Champion, G.D., Laubreaux, L., Nicholas, M.K., Ziegler, J.B. & McInerney, M. (1997). An investigation of the placebo effect and age-related factors in the report of needle pain from venipuncture in children. *Pain* 72 (3) 383-391.
- [31] Kemeny, M.E., Rosenwasser, L.J., Panettieri, R.A., Rose, R.M., Berg-Smith, S.M. & Kline, J.N. (2007). Placebo response in asthma: A robust and objective phenomenon. *Journal of Allergy and Clinical Immunology* 119 (6) 1375-1381.
- [32] Knipschild, P. & Arntz, A. (2005). Pain patients in a randomized trial did not show a significant effect of a positive consultation. *Journal of Clinical Epidemiology* 58 (7) 708-713.
- [33] Lauder, G.R., McQuillan, P.J. & Pickering, R.M. (1995). Psychological adjunct to perioperative antiemesis. *British Journal of Anaesthesia* 74 (3) 266-270.
- [34] Olsson, B., Olsson, B. & Tibblin, G. (1989). Effect of patients' expectations on recovery from acute tonsillitis. *Family Practice* 6 (3) 188-192.
- [35] Petersen, G.L., Finnerup, N.B., Grosen, K., Pilegaard, H.K., Tracey, I., Benedetti, F., Price, D.D., Jensen, T.S. & Vase, L. (2014). Expectations and positive emotional feelings accompany reductions in ongoing and evoked neuropathic pain following placebo interventions. *Pain* 155 (12) 2687-2698.
- [36] Petersen, G.L., Finnerup, N.B., Nørskov, K.N., Grosen, K., Pilegaard, H.K., Benedetti, F., Price, D.D., Jensen, T.S. & Vase, L. (2012). Placebo manipulations reduce hyperalgesia in neuropathic pain. *Pain* 153 (6) 1292-1300.
- [37] Phillips, R.J. (2006). An Examination of the Impact of Self-efficacy Interventions versus Outcome Expectancy Interventions on Efficacy Beliefs, Expectancy Beliefs, Participation, and Functional Performance in a Population of Medical Rehabilitation Patients: Fielding Graduate University. Available at: <https://search.proquest.com/openview/7d7efb8d0f3b971708f30b77b22c1563/1?pq-origsite=gscholar&cbl=18750&diss=y>
- [38] Resnick, B.M. (1996). Self-efficacy in Geriatric Rehabilitation. University of Maryland. Available at: <https://archive.hshsl.umaryland.edu/handle/10713/1478>
- [39] Ronel, J., Mehilli, J., Ladwig, K.H., Blättler, H., Oversohl, N., Byrne, R.A., Bauer, A., Schneider, S., Linde, K., Henningsen, P., Lahmann, C., Noll-Hussong, M. & Meissner, K. (2011). Effects of verbal suggestion on coronary arteries: results of a randomized controlled experimental investigation during coronary angiography. *American Heart Journal* 162 (3) 507-511.
- [40] Suarez-Almazor, M.E., Looney, C., Liu, Y., Cox, V., Pietz, K., Marcus, D.M. & Street, R.L. Jr. (2010). A randomized controlled trial of acupuncture for osteoarthritis of the knee: Effects of patient-provider

- communication. *Arthritis Care & Research* 62 (9) 1229-1236.
- [41] Varelmann, D., Pancaro, C., Cappiello, E.C. & Camann, W.R. (2010). Nocebo-induced hyperalgesia during local anesthetic injection. *Anesthesia & Analgesia* 110 (3) 868-870.
- [42] Wang, F., Shen, X., Xu, S., Liu, Y., Ma, L., Zhao, Q., Fu, D., Pan, Q., Feng, S. & Li, X. (2008). Negative words on surgical wards result in therapeutic failure of patient-controlled analgesia and further release of cortisol after abdominal surgeries. *Minerva Anestesiologica* 74 (7-8) 353-365.
- [43] Szilagyi, A.K., Dioszeghy, C., Benczur, L. & Varga, K. (2007). Effectiveness of psychological support based on positive suggestion with the ventilated patient. *European Journal of Mental Health* 2, 149-170.
- [44] Varga, K., Dioszeghy, C. & Frituz, G. (2007). Suggestive communication with the ventilated patient. *European Journal of Mental Health* 2, 137-147.
- [45] Benedetti, F., Pollo, A., Lopiano, L., Lanotte, M., Vighetti, S. & Rainero, I. (2003). Conscious expectation and unconscious conditioning in analgesic, motor, and hormonal placebo/nocebo responses. [Study 1: Postoperative pain.]. *Journal of Neuroscience* 23 (10) 4315-4323.
- [46] Moerman, D.E. (2002). Meaning, medicine, and the "placebo effect". Cambridge: Cambridge University Press.
- [47] de Craen, A.J., Roos, P.J., Leonard de Vries, A. & Kleijnen, J. (1996). Effect of colour of drugs: systematic review of perceived effect of drugs and of their effectiveness. *British Medical Journal* 313 (7072) 1624-1626.
- [48] Di Blasi, Z., Harkness, E., Ernst, E., Georgiou, A. & Kleijnen, J. (2001). Influence of context effects on health outcomes: a systematic review. *Lancet* 357 (9258) 757-762.
- [49] Benedetti, F., Carlino, E. & Pollo, A. (2011). How placebos change the patient's brain. *Neuropsychopharmacology* 36 (1) 339-354.
- [50] Benedetti, F. (2007). Placebo and endogenous mechanisms of analgesia. *Handbook of Experimental Pharmacology* (177) 393-413.
- [51] Haanstra, T.M., van den Berg, T., Ostelo, R.W., Poolman, R.W., Jansma, I.P., Cuijpers, P. & de Vet, H.C. (2012). Systematic review: Do patient expectations influence treatment outcomes in total knee and total hip arthroplasty? *Health and Quality of Life Outcomes* 10, 152.
- [52] Neuprez, A., Delcour, J.P., Fatemi, F., Gillet, P., Crielaard, J.M., Bruyère, O. & Reginster, J.Y. (2016). Patients' Expectations Impact Their Satisfaction following Total Hip or Knee Arthroplasty. *PLoS One* 11 (12) e0167911.
- [53] Rozanski, A., Bavishi, C., Kubzansky, L.D. & Cohen, R. (2019). Association of Optimism With Cardiovascular Events and All-Cause Mortality: A Systematic Review and Meta-analysis. *JAMA Network Open* 2 (9) e1912200.
- [54] Peerdeman, K.J., van Laarhoven, A.M., Peters, M.L. & Evers, A.W.M. (2016). An Integrative Review of the Influence of Expectancies on Pain. *Frontiers in Psychology* 7, 1270.
- [55] Petty, R.E. & Wegener, D.T. (1998). Attitude change: Multiple roles for persuasion variables. In: *The Handbook of Social Psychology*. Gilbert, D.T., Fiske, S.T. & Lindzey, G. (Eds.), pp 323-390. London: McGraw-Hill.
- [56] Petty, R.E. & Cacioppo, J. (1986). Communication and persuasion: Central and peripheral routes to attitude change. New York: Springer-Verlag.
- [57] Leventhal, H., Meyer, D. & Nerenz, D. (1980). The common sense representation of illness danger. In: *Medical Psychology Vol II*. Rachman, S., (Ed.), pp.7-30. Oxford: Pergamon Press.
- [58] Leventhal, H. & Diefenbach, M. (1991). The active side of illness cognition. In: *Mental representation in health and illness*. Skelton, J.A. & Croyle, R.T. (Eds.), pp. 247-272. New York: Springer Verlag.
- [59] Howe, L.C., Leibowitz, K.A. & Crum, A.J. (2019). When Your Doctor "Gets It" and "Gets You": The Critical Role of Competence and Warmth in the Patient-Provider Interaction. *Frontiers in Psychiatry* 10, 475.
- [60] Ryan, R.M. & Deci, E.L. (2000). Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *American Psychologist* 55 (1) 68-78.
- [61] Bandura, A. (1977). Self-efficacy: toward a unifying theory of behavioral change. *Psychological Review* 84 (2) 191-215.
- [62] Horne, R., Chapman, S.C.E., Parham, R., Freemantle, N., Forbes, A. & Cooper, V. (2013). Understanding Patients' Adherence-Related Beliefs about Medicines Prescribed for Long-Term Conditions: A Meta-Analytic Review of the Necessity-Concerns Framework. *PLoS One* 8 (12) e80633.
- [63] Benedetti, F., Pollo, A., Lopiano, L., Lanotte, M., Vighetti, S. & Rainero, I. (2003). Conscious expectation and unconscious conditioning in analgesic, motor, and hormonal placebo/nocebo responses. [Study 2: State Anxiety]. *Journal of Neuroscience* 23 (10) 4315-4323.
- [64] Benedetti, F., Pollo, A., Lopiano, L., Lanotte, M., Vighetti, S. & Rainero, I. (2003). Conscious expectation and unconscious conditioning in analgesic, motor, and hormonal placebo/nocebo responses. [Study 3: Parkinson's Disease]. *Journal of Neuroscience* 23 (10) 4315-4323.